FORM E

MEDICATIONS & SUPPLEMENTS TO AVOID



Your safety in surgery requires that you disclose all medications, vitamins and supplements that you regularly take. In the TWO WEEKS PRIOR to surgery, you will be required to <u>stop taking</u> certain medications, vitamins and supplements, both those you regularly take, and those that may be taken incidentally for pain or other symptoms.

Please <u>notify</u> Dr. Brad Strawn, or other Sculturra team member, of any and all medications you take during the TWO WEEKS prior to surgery. If you have taken a medication that may put you at risk, it may require that your surgery be rescheduled, or postponed. This is for your safety.

Before you stop taking any prescription drugs, you must receive clearance from the prescribing physician. Please notify our office immediately if you do not receive clearance to stop taking your prescribed medications.

The following drugs could cause life-threatening problems with surgery. If you are on these you must discuss this with the doctor. Patients who take these drugs may require laboratory tests and a consultation with their physician to determine when they may safely undergo a surgical procedure.

Coumadin Elmiron-IC
Methotrexate Plavix
Fragmin and other Low Molecular Weight Heparin Drugs Persantine

Aspirin and aspirin-containing medications and anti-inflammatory agents must not be taken in the TWO WEEKS prior to surgery. Always read the active ingredients on any over the counter or prescription drug packaging. Attached is a listing of common drugs containing aspirin for your reference.

All Herbal or Dietary Supplements should be stopped TWO WEEKS prior to surgery. This includes vitamins, fish oil, anti-oxidants supplements, as well as consumption of any form of Green Tea.

There are several categories of additional medications that must not be taken in the TWO WEEKS prior to surgery. A listing of these drugs is attached.

Authorization

- ➤ I have disclosed all of the medications, supplements and herbal remedies I take on a regular or incidental basis to Brad Strawn, M.D., and/or another Scultura team member.
- I understand that I am required to refrain from taking any of the above and below listed medications in the TWO WEEKS prior to surgery. Should I take any of the medications, supplements or herbal remedies I am instructed to avoid, it is my obligation to notify Dr.Strawn. I fully understand that my surgery may have to be rescheduled or postponed in the event that I have not complied with these medication restrictions listed.
- I also understand that it is my responsibility to obtain clearance from the prescribing physician before I stop taking any of my regularly prescribed medications.

Patient Signature or Personal Representative	Date
Printed Name of Patient or Personal Representative	Relationship of Personal Representative to the Patient
	Signature of Practice Representative and Witness
	orginatare of Fraction Representative and Witness

FORM E

MEDICATIONS & SUPPLEMENTS TO AVOID



SUMMARY OF MEDICATIONS, VITAMINS AND SUPPLEMENTS TO AVOID

Aspirin and aspirin containing medications, include, but are not limited to the following:

Alka Seltzer Carisprodal Compound Momentum
Alka Selzer Plus Cope Norgesic
Anacin Darvon Compound Norgesic Forte
APAC Tablets Darvon Compound 65 Orphagesic
APC Tablets Doan's Pills Orphengesic Forte

Arthritis Pain Reliever Ecotrin PAC
Arthropan Emprin Compound Pamparin
Aspirin Tablets USP Emprin Compuund #3 Percodan
ASA Encaprin Percodan Demi
ASA + Codeine Equagesic Propox Compound

Ascriptin Excedrin Robixisal Asperbuf Fiorinal Sine Off

Aspergum Fiorinal#3 Sodium Salicylate
Axotoal Lortab ASA SOMA Compound
BAC Magnaprin Synalgos

BACMagnaprinSynalgosBayer AspirinMeasurinSynalgosBufferinMediprenTrigesicButalbital ComoundMidolVanquish

Anti-inflammatory medications include but are not limited to the following:

AdvilFeldeneNaproxenAleveIbuprofenNuprinAnaproxIfenOrudisAnsaidIndocinOvuvail

Butazoladin Indomethocin Phenylbutazone

CataflamMeclomenRutenClinotilMotrinTolectinDayproNalfonToradolDolobidNaprosynVoltaren

Additional medications to avoid, include, but are not limited to the following:

St. John's Wort Chloratrimeton Multi-vitamin Mysteclin f Surmontil Clinoril Nicobid Elavil **Tagamet** Oraflex Tenuate Dospan Endep Tetracycline Etiafon Pamelor Triavil Flagyl Parnate Flexoril Phendimetrazine Vibramycin Vitamin E **Imitrex** Phentermine Zomax Lioresal Ru-Tuss